

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
1 July 2004 (01.07.2004)

PCT

(10) International Publication Number  
WO 2004/054351 A2

- (51) International Patent Classification<sup>7</sup>: **A01H**
- (21) International Application Number:  
PCT/US2003/041146
- (22) International Filing Date:  
18 December 2003 (18.12.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/434,763 18 December 2002 (18.12.2002) US
- (71) Applicant (for all designated States except US): **AGRI-  
NOMICS LLC** [US/US]; 16160 S.W. Upper Boones Ferry  
Road, Portland, OR 97224-7744 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LIGHTNER,  
Jonathan** [US/US]; 28750 S. Heisinger Lane, Mulino, OR  
97042 (US). **CLENDENNEN, Stephanie, K.** [US/US];  
P.O. Box 6332, Kingsport, TN 37663 (US).
- (74) Agents: **BRUNELLE, Jan, P.** et al.; Exelixis, Inc., 170  
Harbor Way, P.O. Box 511, South San Francisco, CA  
94083-0511 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,  
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,  
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,  
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished  
upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: GENERATION OF PLANTS WITH ALTERED OIL CONTENT

(57) Abstract: The present invention is directed to plants that display an altered oil content phenotype due to altered expression of a HIO30 nucleic acid. The invention is further directed to methods of generating plants with an altered oil content phenotype.

WO 2004/054351 A2

## GENERATION OF PLANTS WITH ALTERED OIL CONTENT

### REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. provisional patent application no.

5 60/434,763 filed 12/18/2002, the contents of which are hereby incorporated by reference

### BACKGROUND OF THE INVENTION

The ability to manipulate the composition of crop seeds, particularly the content and composition of seed oils, has important applications in the agricultural industries, relating both to processed food oils and to oils for animal feeding. Seeds of agricultural crops contain a variety of valuable constituents, including oil, protein and starch. Industrial processing can separate some or all of these constituents for individual sale in specific applications. For instance, nearly 60% of the US soybean crop is crushed by the soy processing industry. Soy processing yields purified oil, which is sold at high value, while the remainder is sold principally for lower value livestock feed (US Soybean Board, 2001 Soy Stats). Canola seed is crushed to produce oil and the co-product canola meal (Canola Council of Canada). Nearly 20% of the 1999/2000 US corn crop was industrially refined, primarily for production of starch, ethanol and oil (Corn Refiners Association). Thus, it is often desirable to maximize oil content of seeds. For instance, for processed oilseeds such as soy and canola, increasing the absolute oil content of the seed will increase the value of such grains. For processed corn it may be desired to either increase or decrease oil content, depending on utilization of other major constituents. Decreasing oil may improve the quality of isolated starch by reducing undesired flavors associated with oil oxidation. Alternatively, in ethanol production, where flavor is unimportant, increasing oil content may increase overall value. In many fed grains, such as corn and wheat, it is desirable to increase seed oil content, because oil has higher energy content than other seed constituents such as carbohydrate. Oilseed processing, like most grain processing businesses, is a capital-intensive business; thus small shifts in the distribution of products from the low valued components to the high value oil component can have substantial economic impacts for grain processors.

Biotechnological manipulation of oils can provide compositional alteration and improvement of oil yield. Compositional alterations include high oleic soybean and corn oil (US Pat Nos 6,229,033 and 6,248,939), and laurate-containing seeds (US Pat No 5,639,790), among others. Work in compositional alteration has predominantly focused

on processed oilseeds but has been readily extendable to non-oilseed crops, including corn. While there is considerable interest in increasing oil content, the only currently practiced biotechnology in this area is High-Oil Corn (HOC) technology (DuPont, US Pat. No. 5,704,160). HOC employs high oil pollinators developed by classical selection  
5 breeding along with elite (male-sterile) hybrid females in a production system referred to as TopCross. The TopCross High Oil system raises harvested grain oil content in maize from ~3.5% to ~7%, improving the energy content of the grain.

While it has been fruitful, the HOC production system has inherent limitations. First, the system of having a low percentage of pollinators responsible for an entire field's  
10 seed set contains inherent risks, particularly in drought years. Second, oil contents in current HOC fields have plateaued at about 9% oil. Finally, high-oil corn is not primarily a biochemical change, but rather an anatomical mutant (increased embryo size) that has the indirect result of increasing oil content. For these reasons, an alternative high oil strategy, particularly one that derives from an altered biochemical output, would be  
15 especially valuable.

The most obvious target crops for the processed oil market are soy and rapeseed, and a large body of commercial work (e.g., US Pat No: 5,952,544; PCT application WO9411516) demonstrates that *Arabidopsis* is an excellent model for oil metabolism in these crops. Biochemical screens of seed oil composition have identified *Arabidopsis*  
20 genes for many critical biosynthetic enzymes and have led to identification of agronomically important gene orthologs. For instance, screens using chemically mutagenized populations have identified lipid mutants whose seeds display altered fatty acid composition (Lemieux B, et al. 1990, Theor Appl Genet 80, 234-240; James DW and Dooner HK (1990) Theor Appl Genet 80, 241-245). T-DNA mutagenesis screens  
25 (Feldmann *et al.*, *Science* 243: 1351-1354, 1989) that detected altered fatty acid composition identified the omega 3 desaturase (*FAD3*) and delta-12 desaturase (*FAD2*) genes (US Pat No 5952544; Yadav NS et al. (1993) Plant Physiol 103, 467-476; Okuley et al., Plant Cell. 1994 Jan;6(1):147-58). A screen which focused on oil content rather than oil quality, analyzed chemically-induced mutants for wrinkled seeds or altered seed  
30 density, from which altered seed oil content was inferred (Focks N and Benning C, Plant Physiol 118:91-101, 1998). Another screen, designed to identify enzymes involved in production of very long chain fatty acids, identified a mutation in the gene encoding a diacylglycerol acyltransferase (DGAT) as being responsible for reduced triacyl glycerol accumulation in seeds (Katavic V et al, Plant Physiol. 1995 May;108(1):399-409). It was

further shown that seed-specific over-expression of the DGAT cDNA was associated with increased seed oil content (Jako et al., Plant Physiol. 2001 Jun;126(2):861-74).

Activation tagging in plants refers to a method of generating random mutations by insertion of a heterologous nucleic acid construct comprising regulatory sequences (e.g., an enhancer) into a plant genome. The regulatory sequences can act to enhance transcription of one or more native plant genes; accordingly, activation tagging is a fruitful method for generating gain-of-function, generally dominant mutants (see, e.g., Hayashi et al., Science (1992) 258: 1350-1353; Weigel et al., Plant Physiology (2000) 122:1003-1013). The inserted construct provides a molecular tag for rapid identification of the native plant whose mis-expression causes the mutant phenotype. Activation tagging may also cause loss-of-function phenotypes. The insertion may result in disruption of a native plant gene, in which case the phenotype is generally recessive.

Activation tagging has been used in various species, including tobacco and *Arabidopsis*, to identify many different kinds of mutant phenotypes and the genes associated with these phenotypes (Wilson et al., Plant Cell (1996) 8:659-671, Schaffer et al., Cell (1998) 93: 1219-1229; Fridborg et al., Plant Cell (1999) 11: 1019-1032; Kardailsky et al., Science (1999) 286:1962-1965); Christensen S et al., 9<sup>th</sup> International Conference on *Arabidopsis* Research. Univ. of Wisconsin-Madison, June 24-28, 1998. Abstract 165).

## SUMMARY OF THE INVENTION

The invention provides a transgenic plant having a high oil phenotype. The transgenic plant comprises a transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO30 polypeptide. In preferred embodiments, the transgenic plant is selected from the group consisting of rapeseed, soy, corn, sunflower, cotton, cocoa, safflower, oil palm, coconut palm, flax, castor and peanut. The invention further provides a method of producing oil comprising growing the transgenic plant and recovering oil from said plant.

The transgenic plant of the invention is produced by a method that comprises introducing into progenitor cells of the plant a plant transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO30 polypeptide, and growing the transformed progenitor cells to produce a transgenic plant, wherein the HIO30 polynucleotide sequence is expressed causing the high oil phenotype.

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions

Unless otherwise indicated, all technical and scientific terms used herein have the same meaning as they would to one skilled in the art of the present invention.

5 Practitioners are particularly directed to Sambrook *et al.* Molecular Cloning: A Laboratory Manual (Second Edition), Cold Spring Harbor Press, Plainview, N.Y., 1989, and Ausubel *et al.* Current Protocols in Molecular Biology, John Wiley & Sons, New York, N.Y., 1993, for definitions and terms of the art. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may  
10 vary.

As used herein, the term "vector" refers to a nucleic acid construct designed for transfer between different host cells. An "expression vector" refers to a vector that has the ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of  
15 appropriate expression vectors is within the knowledge of those having skill in the art.

A "heterologous" nucleic acid construct or sequence has a portion of the sequence that is not native to the plant cell in which it is expressed. Heterologous, with respect to a control sequence refers to a control sequence (*i.e.* promoter or enhancer) that does not function in nature to regulate the same gene the expression of which it is currently  
20 regulating. Generally, heterologous nucleic acid sequences are not endogenous to the cell or part of the genome in which they are present, and have been added to the cell, by infection, transfection, microinjection, electroporation, or the like. A "heterologous" nucleic acid construct may contain a control sequence/DNA coding sequence combination that is the same as, or different from a control sequence/DNA coding sequence  
25 combination found in the native plant.

As used herein, the term "gene" means the segment of DNA involved in producing a polypeptide chain, which may or may not include regions preceding and following the coding region, *e.g.* 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments  
30 (exons) and non-transcribed regulatory sequence.

As used herein, "recombinant" includes reference to a cell or vector, that has been modified by the introduction of a heterologous nucleic acid sequence or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or

express native genes that are otherwise abnormally expressed, under expressed or not expressed at all as a result of deliberate human intervention.

As used herein, the term "gene expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation; accordingly, "expression" may refer to either a polynucleotide or polypeptide sequence, or both. Sometimes, expression of a polynucleotide sequence will not lead to protein translation. "Over-expression" refers to increased expression of a polynucleotide and/or polypeptide sequence relative to its expression in a wild-type (or other reference [e.g., non-transgenic]) plant and may relate to a naturally-occurring or non-naturally occurring sequence. "Ectopic expression" refers to expression at a time, place, and/or increased level that does not naturally occur in the non-altered or wild-type plant. "Under-expression" refers to decreased expression of a polynucleotide and/or polypeptide sequence, generally of an endogenous gene, relative to its expression in a wild-type plant. The terms "mis-expression" and "altered expression" encompass over-expression, under-expression, and ectopic expression.

The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

As used herein, a "plant cell" refers to any cell derived from a plant, including cells from undifferentiated tissue (e.g., callus) as well as plant seeds, pollen, progagules and embryos.

As used herein, the terms "native" and "wild-type" relative to a given plant trait or phenotype refers to the form in which that trait or phenotype is found in the same variety of plant in nature.

As used herein, the term "modified" regarding a plant trait, refers to a change in the phenotype of a transgenic plant relative to the similar non-transgenic plant. An "altered oil content phenotype" refers to measurable phenotype of a genetically modified plant, where the plant displays a statistically significant increase or decrease in overall oil content (i.e., the percentage of seed mass that is oil), as compared to the similar, but non-modified plant. A high oil phenotype refers to an increase in overall oil content.

As used herein, a "mutant" polynucleotide sequence or gene differs from the corresponding wild type polynucleotide sequence or gene either in terms of sequence or expression, where the difference contributes to a modified plant phenotype or trait. Relative to a plant or plant line, the term "mutant" refers to a plant or plant line which has  
5 a modified plant phenotype or trait, where the modified phenotype or trait is associated with the modified expression of a wild type polynucleotide sequence or gene.

As used herein, the term "T1" refers to the generation of plants from the seed of T0 plants. The T1 generation is the first set of transformed plants that can be selected by application of a selection agent, *e.g.*, an antibiotic or herbicide, for which the transgenic  
10 plant contains the corresponding resistance gene. The term "T2" refers to the generation of plants by self-fertilization of the flowers of T1 plants, previously selected as being transgenic. T3 plants are generated from T2 plants, etc. As used herein, the "direct progeny" of a given plant derives from the seed (or, sometimes, other tissue) of that plant and is in the immediately subsequent generation; for instance, for a given lineage, a T2  
15 plant is the direct progeny of a T1 plant. The "indirect progeny" of a given plant derives from the seed (or other tissue) of the direct progeny of that plant, or from the seed (or other tissue) of subsequent generations in that lineage; for instance, a T3 plant is the indirect progeny of a T1 plant.

As used herein, the term "plant part" includes any plant organ or tissue, including,  
20 without limitation, seeds, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. Plant cells can be obtained from any plant organ or tissue and cultures prepared therefrom. The class of plants which can be used in the methods of the present invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledenous  
25 and dicotyledenous plants.

As used herein, "transgenic plant" includes a plant that comprises within its genome a heterologous polynucleotide. The heterologous polynucleotide can be either stably integrated into the genome, or can be extra-chromosomal. Preferably, the polynucleotide of the present invention is stably integrated into the genome such that the  
30 polynucleotide is passed on to successive generations. A plant cell, tissue, organ, or plant into which the heterologous polynucleotides have been introduced is considered "transformed", "transfected", or "transgenic". Direct and indirect progeny of transformed plants or plant cells that also contain the heterologous polynucleotide are also considered transgenic.

### **Identification of Plants with an Altered Oil Content Phenotype**

We used an *Arabidopsis* activation tagging screen to identify the association between the gene we have designated "HIO30," (At3g52260; GI#22331747:54-938), and an altered oil content phenotype (specifically, a high oil phenotype). Briefly, and as  
5 further described in the Examples, a large number of *Arabidopsis* plants were mutated with the pSKI015 vector, which comprises a T-DNA from the Ti plasmid of *Agrobacterium tumefaciens*, a viral enhancer element, and a selectable marker gene (Weigel *et al*, *supra*). When the T-DNA inserts into the genome of transformed plants, the enhancer element can cause up-regulation genes in the vicinity, generally within about 10  
10 kilobase (kb) of the insertion. T1 plants were exposed to the selective agent in order to specifically recover transformed plants that expressed the selectable marker and therefore harbored T-DNA insertions. Samples of approximately 15-20 T2 seeds were collected from transformed T1 plants, and lipids were extracted from whole seeds. Gas chromatography (GC) analysis was performed to determine fatty acid content and  
15 composition of seed samples.

An *Arabidopsis* line that showed a high-oil phenotype was identified, wherein oils (i.e., fatty acids) constituted approximately 35% of seed mass. The association of the HIO30 gene with the high oil phenotype was discovered by analysis of the genomic DNA sequence flanking the T-DNA insertion in the identified line. Accordingly, HIO30 genes  
20 and/or polypeptides may be employed in the development of genetically modified plants having a modified oil content phenotype ("a HIO30 phenotype"). HIO30 genes may be used in the generation of oilseed crops that provide improved oil yield from oilseed processing and in the generation of feed grain crops that provide increased energy for animal feeding. HIO30 genes may further be used to increase the oil content of specialty  
25 oil crops, in order to augment yield of desired unusual fatty acids. Transgenic plants that have been genetically modified to express HIO30 can be used in the production of oil, wherein the transgenic plants are grown, and oil is obtained from plant parts (e.g. seed) using standard methods.

### **HIO30 Nucleic Acids and Polypeptides**

*Arabidopsis* HIO30 nucleic acid (genomic DNA) sequence is provided in SEQ ID NO:1 and in Genbank entry GI#22331747:54-938. The corresponding protein sequence is provided in SEQ ID NO:2 and in GI#22331748. Nucleic acids and/or proteins that are orthologs or paralogs of *Arabidopsis* HIO30, are described in Example 3 below.



As used herein, the term "HIO30 polypeptide" refers to a full-length HIO30 protein or a fragment, derivative (variant), or ortholog thereof that is "functionally active," meaning that the protein fragment, derivative, or ortholog exhibits one or more of the functional activities associated with the polypeptide of SEQ ID NO:2. In one preferred embodiment, a functionally active HIO30 polypeptide causes an altered oil content phenotype when mis-expressed in a plant. In a further preferred embodiment, mis-expression of the HIO30 polypeptide causes a high oil phenotype in a plant. In another embodiment, a functionally active HIO30 polypeptide is capable of rescuing defective (including deficient) endogenous HIO30 activity when expressed in a plant or in plant cells; the rescuing polypeptide may be from the same or from a different species as that with defective activity. In another embodiment, a functionally active fragment of a full length HIO30 polypeptide (i.e., a native polypeptide having the sequence of SEQ ID NO:2 or a naturally occurring ortholog thereof) retains one of more of the biological properties associated with the full-length HIO30 polypeptide, such as signaling activity, binding activity, catalytic activity, or cellular or extra-cellular localizing activity. A HIO30 fragment preferably comprises a HIO30 domain, such as a C- or N-terminal or catalytic domain, among others, and preferably comprises at least 10, preferably at least 20, more preferably at least 25, and most preferably at least 50 contiguous amino acids of a HIO30 protein. Functional domains can be identified using the PFAM program (Bateman A et al., 1999 Nucleic Acids Res 27:260-262; website at pfam.wustl.edu). A preferred HIO30 fragment comprises a pseudouridylate synthase domain (PFam00849). Functionally active variants of full-length HIO30 polypeptides or fragments thereof include polypeptides with amino acid insertions, deletions, or substitutions that retain one of more of the biological properties associated with the full-length HIO30 polypeptide. In some cases, variants are generated that change the post-translational processing of a HIO30 polypeptide. For instance, variants may have altered protein transport or protein localization characteristics or altered protein half-life compared to the native polypeptide.

As used herein, the term "HIO30 nucleic acid" encompasses nucleic acids with the sequence provided in or complementary to the sequence provided in SEQ ID NO:1, as well as functionally active fragments, derivatives, or orthologs thereof. A HIO30 nucleic acid of this invention may be DNA, derived from genomic DNA or cDNA, or RNA.

In one embodiment, a functionally active HIO30 nucleic acid encodes or is complementary to a nucleic acid that encodes a functionally active HIO30 polypeptide. Included within this definition is genomic DNA that serves as a template for a primary

RNA transcript (i.e., an mRNA precursor) that requires processing, such as splicing, before encoding the functionally active HIO30 polypeptide. A HIO30 nucleic acid can include other non-coding sequences, which may or may not be transcribed; such sequences include 5' and 3' UTRs, polyadenylation signals and regulatory sequences that control gene expression, among others, as are known in the art. Some polypeptides require processing events, such as proteolytic cleavage, covalent modification, etc., in order to become fully active. Accordingly, functionally active nucleic acids may encode the mature or the pre-processed HIO30 polypeptide, or an intermediate form. A HIO30 polynucleotide can also include heterologous coding sequences, for example, sequences that encode a marker included to facilitate the purification of the fused polypeptide, or a transformation marker.

In another embodiment, a functionally active HIO30 nucleic acid is capable of being used in the generation of loss-of-function HIO30 phenotypes, for instance, via antisense suppression, co-suppression, etc.

In one preferred embodiment, a HIO30 nucleic acid used in the methods of this invention comprises a nucleic acid sequence that encodes or is complementary to a sequence that encodes a HIO30 polypeptide having at least 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95% or more sequence identity to the polypeptide sequence presented in SEQ ID NO:2.

In another embodiment a HIO30 polypeptide of the invention comprises a polypeptide sequence with at least 50% or 60% identity to the HIO30 polypeptide sequence of SEQ ID NO:2, and may have at least 70%, 80%, 85%, 90% or 95% or more sequence identity to the HIO30 polypeptide sequence of SEQ ID NO:2. In another embodiment, a HIO30 polypeptide comprises a polypeptide sequence with at least 50%, 60%, 70%, 80%, 85%, 90% or 95% or more sequence identity to a functionally active fragment of the polypeptide presented in SEQ ID NO:2, such as a pseudouridylate synthase domain. In yet another embodiment, a HIO30 polypeptide comprises a polypeptide sequence with at least 50%, 60 %, 70%, 80%, or 90% identity to the polypeptide sequence of SEQ ID NO:2 over its entire length and comprises a pseudouridylate synthase domain.

In another aspect, a HIO30 polynucleotide sequence is at least 50% to 60% identical over its entire length to the HIO30 nucleic acid sequence presented as SEQ ID NO:1, or nucleic acid sequences that are complementary to such a HIO30 sequence, and may comprise at least 70%, 80%, 85%, 90% or 95% or more sequence identity to the

HIO30 sequence presented as SEQ ID NO:1 or a functionally active fragment thereof, or complementary sequences.

As used herein, "percent (%) sequence identity" with respect to a specified subject sequence, or a specified portion thereof, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410; website at [blast.wustl.edu/blast/README.html](http://blast.wustl.edu/blast/README.html)) with search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A "% identity value" is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation. A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that selectively hybridize to the nucleic acid sequence of SEQ ID NO:1. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are well known (see, *e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing

the nucleotide sequence of SEQ ID NO:1 under stringent hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1 h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate). In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS. Alternatively, low stringency conditions can be used that comprise: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

As a result of the degeneracy of the genetic code, a number of polynucleotide sequences encoding a HIO30 polypeptide can be produced. For example, codons may be selected to increase the rate at which expression of the polypeptide occurs in a particular host species, in accordance with the optimum codon usage dictated by the particular host organism (see, e.g., Nakamura et al, 1999, Nucleic Acids Res 27:292). Such sequence variants may be used in the methods of this invention.

The methods of the invention may use orthologs of the *Arabidopsis* HIO30. Methods of identifying the orthologs in other plant species are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *Arabidopsis*, may correspond to multiple genes (paralogs) in another. As used herein, the term "orthologs" encompasses paralogs. When sequence data is available for a particular plant species, orthologs are generally identified by sequence homology analysis, such as

BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210).

5 Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD *et al.*, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the

10 tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. Nucleic acid hybridization methods may also be used to find orthologous genes and are preferred when sequence data are not available. Degenerate PCR and screening of cDNA or genomic DNA libraries are

15 common methods for finding related gene sequences and are well known in the art (see, e.g., Sambrook, *supra*; Dieffenbach C and Dveksler G (Eds.) PCR Primer: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY, 1989). For instance, methods for generating a cDNA library from the plant species of interest and probing the library with partially homologous gene probes are described in Sambrook *et al.*, *supra*. A highly

20 conserved portion of the *Arabidopsis* HIO30 coding sequence may be used as a probe. HIO30 ortholog nucleic acids may hybridize to the nucleic acid of SEQ ID NO:1 under high, moderate, or low stringency conditions. After amplification or isolation of a segment of a putative ortholog, that segment may be cloned and sequenced by standard techniques and utilized as a probe to isolate a complete cDNA or genomic clone.

25 Alternatively, it is possible to initiate an EST project to generate a database of sequence information for the plant species of interest. In another approach, antibodies that specifically bind known HIO30 polypeptides are used for ortholog isolation (see, e.g., Harlow E and Lane D, Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, 1999, New York). Western blot analysis can determine that a HIO30

30 ortholog (i.e., an orthologous protein) is present in a crude extract of a particular plant species. When reactivity is observed, the sequence encoding the candidate ortholog may be isolated by screening expression libraries representing the particular plant species. Expression libraries can be constructed in a variety of commercially available vectors, including lambda gt11, as described in Sambrook, *et al.*, *supra*. Once the candidate

ortholog(s) are identified by any of these means, candidate orthologous sequence are used as bait (the "query") for the reverse BLAST against sequences from *Arabidopsis* or other species in which HIO30 nucleic acid and/or polypeptide sequences have been identified.

HIO30 nucleic acids and polypeptides may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR), as previously described, are well known in the art. Alternatively, nucleic acid sequence may be synthesized. Any known method, such as site directed mutagenesis (Kunkel TA *et al.*, *Methods Enzymol.* 204:125-39, 1991), may be used to introduce desired changes into a cloned nucleic acid.

In general, the methods of the invention involve incorporating the desired form of the HIO30 nucleic acid into a plant expression vector for transformation of in plant cells, and the HIO30 polypeptide is expressed in the host plant.

An isolated HIO30 nucleic acid molecule is other than in the form or setting in which it is found in nature and is identified and separated from least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the HIO30 nucleic acid. However, an isolated HIO30 nucleic acid molecule includes HIO30 nucleic acid molecules contained in cells that ordinarily express HIO30 where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

#### **Generation of Genetically Modified Plants with an Altered Oil Content Phenotype**

HIO30 nucleic acids and polypeptides may be used in the generation of genetically modified plants having a modified oil content phenotype. As used herein, a "modified oil content phenotype" may refer to modified oil content in any part of the plant; the modified oil content is often observed in seeds. In a preferred embodiment, altered expression of the HIO30 gene in a plant is used to generate plants with a high oil phenotype.

The methods described herein are generally applicable to all plants. Although activation tagging and gene identification is carried out in *Arabidopsis*, the HIO30 gene (or an ortholog, variant or fragment thereof) may be expressed in any type of plant. In a preferred embodiment, the invention is directed to oil-producing plants, which produce and store triacylglycerol in specific organs, primarily in seeds. Such species include soybean (*Glycine max*), rapeseed and canola (including *Brassica napus*, *B. campestris*), sunflower (*Helianthus annuus*), cotton (*Gossypium hirsutum*), corn (*Zea mays*), cocoa

(*Theobroma cacao*), safflower (*Carthamus tinctorius*), oil palm (*Elaeis guineensis*), coconut palm (*Cocos nucifera*), flax (*Linum usitatissimum*), castor (*Ricinus communis*) and peanut (*Arachis hypogaea*). The invention may also be directed to fruit- and vegetable-bearing plants, grain-producing plants, nut-producing plants, rapid cycling  
5 *Brassica* species, alfalfa (*Medicago sativa*), tobacco (*Nicotiana*), turfgrass (*Poaceae* family), other forage crops, and wild species that may be a source of unique fatty acids.

The skilled artisan will recognize that a wide variety of transformation techniques exist in the art, and new techniques are continually becoming available. Any technique that is suitable for the target host plant can be employed within the scope of the present  
10 invention. For example, the constructs can be introduced in a variety of forms including, but not limited to as a strand of DNA, in a plasmid, or in an artificial chromosome. The introduction of the constructs into the target plant cells can be accomplished by a variety of techniques, including, but not limited to *Agrobacterium*-mediated transformation, electroporation, microinjection, microprojectile bombardment calcium-phosphate-DNA  
15 co-precipitation or liposome-mediated transformation of a heterologous nucleic acid. The transformation of the plant is preferably permanent, *i.e.* by integration of the introduced expression constructs into the host plant genome, so that the introduced constructs are passed onto successive plant generations. Depending upon the intended use, a heterologous nucleic acid construct comprising an HIO30 polynucleotide may encode the  
20 entire protein or a biologically active portion thereof.

In one embodiment, binary Ti-based vector systems may be used to transfer polynucleotides. Standard *Agrobacterium* binary vectors are known to those of skill in the art, and many are commercially available (e.g., pBI121 Clontech Laboratories, Palo Alto, CA).

25 The optimal procedure for transformation of plants with *Agrobacterium* vectors will vary with the type of plant being transformed. Exemplary methods for *Agrobacterium*-mediated transformation include transformation of explants of hypocotyl, shoot tip, stem or leaf tissue, derived from sterile seedlings and/or plantlets. Such transformed plants may be reproduced sexually, or by cell or tissue culture.  
30 *Agrobacterium* transformation has been previously described for a large number of different types of plants and methods for such transformation may be found in the scientific literature. Of particular relevance are methods to transform commercially important crops, such as rapeseed (De Block et al., Plant Physiol. (1989) 91:694-701), sunflower (Everett et

al., Bio/Technology (1987) 5:1201), and soybean (Christou et al., Proc. Natl. Acad. Sci USA (1989) 86:7500-7504; Kline et al., Nature (1987) 327:70).

Expression (including transcription and translation) of HIO30 may be regulated with respect to the level of expression, the tissue type(s) where expression takes place and/or developmental stage of expression. A number of heterologous regulatory sequences (e.g., promoters and enhancers) are available for controlling the expression of a HIO30 nucleic acid. These include constitutive, inducible and regulatable promoters, as well as promoters and enhancers that control expression in a tissue- or temporal-specific manner. Exemplary constitutive promoters include the raspberry E4 promoter (U.S. Patent Nos. 5,783,393 and 5,783,394), the 35S CaMV (Jones JD *et al.*, Transgenic Res 1:285-297 1992), the CsVMV promoter (Verdaguer B *et al.*, Plant Mol Biol 37:1055-1067, 1998) and the melon actin promoter (published PCT application WO0056863). Exemplary tissue-specific promoters include the tomato E4 and E8 promoters (U.S. Pat. No. 5,859,330) and the tomato 2AII gene promoter (Van Haaren MJJ *et al.*, Plant Mol Bio 21:625-640, 1993).

In one preferred embodiment, HIO30 expression is under control of regulatory sequences from genes whose expression is associated with early seed and/or embryo development. Legume genes whose promoters are associated with early seed and embryo development include *V. faba legumin* (Baumlein et al., 1991, Mol Gen Genet 225:121-8; Baumlein et al., 1992, Plant J 2:233-9), *V. faba usp* (Fiedler et al., 1993, Plant Mol Biol 22:669-79), pea *convicilin* (Bown et al., 1988, Biochem J 251:717-26), pea *lectin* (dePater et al., 1993, Plant Cell 5:877-86), *P. vulgaris beta phaseolin* (Bustos et al., 1991, EMBO J 10:1469-79), *P. vulgaris DLEC2* and *PHS* [beta] (Bobb et al, 1997, Nucleic Acids Res 25:641-7), and soybean *beta-Conglycinin*, 7S storage protein (Chamberland et al., 1992, Plant Mol Biol 19:937-49). Cereal genes whose promoters are associated with early seed and embryo development include rice *glutelin* ("GluA-3," Yoshihara and Takaiwa, 1996, Plant Cell Physiol 37:107-11; "GluB-1," Takaiwa et al., 1996, Plant Mol Biol 30:1207-21; Washida et al., 1999, Plant Mol Biol 40:1-12; "Gt3," Leisy et al., 1990, Plant Mol Biol 14:41-50), rice *prolamin* (Zhou & Fan, 1993, Transgenic Res 2:141-6), wheat *prolamin* (Hammond-Kosack et al., 1993, EMBO J 12:545-54), maize *zein* (Z4, Matzke et al., 1990, Plant Mol Biol 14:323-32), and barley *B-hordeins* (Entwistle et al., 1991, Plant Mol Biol 17:1217-31). Other genes whose promoters are associated with early seed and embryo development include oil palm GLO7A (7S globulin, Morcillo et al., 2001, Physiol Plant 112:233-243), *Brassica napus napin*, 2S storage protein, and napA gene (Josefsson et al.,



1987, J Biol Chem 262:12196-201; Stalberg et al., 1993, Plant Mol Biol 1993 23:671-83; Ellerstrom et al., 1996, Plant Mol Biol 32:1019-27), *Brassica napus oleosin* (Keddie et al., 1994, Plant Mol Biol 24:327-40), *Arabidopsis oleosin* (Plant et al., 1994, Plant Mol Biol 25:193-205), *Arabidopsis* FAE1 (Rossak et al., 2001, Plant Mol Biol 46:717-25),  
5 *Canavalia gladiata* conA (Yamamoto et al., 1995, Plant Mol Biol 27:729-41), and *Catharanthus roseus* strictosidine synthase (Str, Ouwerkerk and Memelink, 1999, Mol Gen Genet 261:635-43). In another preferred embodiment, regulatory sequences from genes expressed during oil biosynthesis are used (see, e.g., US Pat No: 5,952, 544). Alternative promoters are from plant storage protein genes (Bevan et al, 1993, Philos  
10 Trans R Soc Lond B Biol Sci 342:209-15).

In yet another aspect, in some cases it may be desirable to inhibit the expression of endogenous HIO30 in a host cell. Exemplary methods for practicing this aspect of the invention include, but are not limited to antisense suppression (Smith, *et al.*, *Nature* 334:724-726, 1988; van der Krol et al., *Biotechniques* (1988) 6:958-976); co-suppression  
15 (Napoli, *et al.*, *Plant Cell* 2:279-289, 1990); ribozymes (PCT Publication WO 97/10328); and combinations of sense and antisense (Waterhouse, *et al.*, *Proc. Natl. Acad. Sci. USA* 95:13959-13964, 1998). Methods for the suppression of endogenous sequences in a host cell typically employ the transcription or transcription and translation of at least a portion of the sequence to be suppressed. Such sequences may be homologous to coding as well  
20 as non-coding regions of the endogenous sequence. Antisense inhibition may use the entire cDNA sequence (Sheehy et al., *Proc. Natl. Acad. Sci. USA* (1988) 85:8805-8809), a partial cDNA sequence including fragments of 5' coding sequence, (Cannon et al., *Plant Molec. Biol.* (1990) 15:39-47), or 3' non-coding sequences (Ch'ng et al., *Proc. Natl. Acad. Sci. USA* (1989) 86:10006-10010). Cosuppression techniques may use the entire cDNA  
25 sequence (Napoli et al., *supra*; van der Krol et al., *The Plant Cell* (1990) 2:291-299) or a partial cDNA sequence (Smith et al., *Mol. Gen. Genetics* (1990) 224:477-481).

Standard molecular and genetic tests may be performed to further analyze the association between a gene and an observed phenotype. Exemplary techniques are  
30 described below.

#### 1. DNA/RNA analysis

The stage- and tissue-specific gene expression patterns in mutant versus wild-type lines may be determined, for instance, by in situ hybridization. Analysis of the methylation status of the gene, especially flanking regulatory regions, may be performed. Other

suitable techniques include overexpression, ectopic expression, expression in other plant species and gene knock-out (reverse genetics, targeted knock-out, viral induced gene silencing [VIGS, see Baulcombe D, *Arch Virol Suppl* 15:189-201, 1999]).

In a preferred application expression profiling, generally by microarray analysis, is used to simultaneously measure differences or induced changes in the expression of many different genes. Techniques for microarray analysis are well known in the art (Schena M *et al.*, *Science* (1995) 270:467-470; Baldwin D *et al.*, *Cur Opin Plant Biol.* 2(2):96-103, 1999; Dangond F, *Physiol Genomics* (2000) 2:53-58; van Hal NL *et al.*, *J Biotechnol* (2000) 78:271-280; Richmond T and Somerville S, *Curr Opin Plant Biol* (2000) 3:108-116). Expression profiling of individual tagged lines may be performed. Such analysis can identify other genes that are coordinately regulated as a consequence of the overexpression of the gene of interest, which may help to place an unknown gene in a particular pathway.

## 2. Gene Product Analysis

Analysis of gene products may include recombinant protein expression, antisera production, immunolocalization, biochemical assays for catalytic or other activity, analysis of phosphorylation status, and analysis of interaction with other proteins via yeast two-hybrid assays.

## 3. Pathway Analysis

Pathway analysis may include placing a gene or gene product within a particular biochemical, metabolic or signaling pathway based on its mis-expression phenotype or by sequence homology with related genes. Alternatively, analysis may comprise genetic crosses with wild-type lines and other mutant lines (creating double mutants) to order the gene in a pathway, or determining the effect of a mutation on expression of downstream "reporter" genes in a pathway.

## Generation of Mutated Plants with an Altered Oil Content Phenotype

The invention further provides a method of identifying plants that have mutations in endogenous HIO30 that confer altered oil content, and generating altered oil content progeny of these plants that are not genetically modified. In one method, called "TILLING" (for targeting induced local lesions in genomes), mutations are induced in the seed of a plant of interest, for example, using EMS treatment. The resulting plants are grown and self-fertilized, and the progeny are used to prepare DNA samples. HIO30-specific PCR is used to identify whether a mutated plant has a HIO30 mutation. Plants

having HIO30 mutations may then be tested for altered oil content, or alternatively, plants may be tested for altered oil content, and then HIO30-specific PCR is used to determine whether a plant having altered oil content has a mutated HIO30 gene. TILLING can identify mutations that may alter the expression of specific genes or the activity of proteins encoded by these genes (see Colbert et al (2001) Plant Physiol 126:480-484; McCallum et al (2000) Nature Biotechnology 18:455-457).

In another method, a candidate gene/Quantitative Trait Locus (QTLs) approach can be used in a marker-assisted breeding program to identify alleles of or mutations in the HIO30 gene or orthologs of HIO30 that may confer altered oil content (see Bert et al., Theor Appl Genet. 2003 Jun;107(1):181-9; and Lionneton et al, Genome. 2002 Dec;45(6):1203-15). Thus, in a further aspect of the invention, a HIO30 nucleic acid is used to identify whether a plant having altered oil content has a mutation in endogenous HIO30 or has a particular allele that causes altered oil content.

All publications cited herein are expressly incorporated herein by reference for the purpose of describing and disclosing compositions and methodologies that might be used in connection with the invention. All cited patents, patent applications, and sequence information in referenced websites and public databases are also incorporated by reference.

## EXAMPLES

### EXAMPLE 1

#### Generation of Plants with a HIO30 Phenotype by Transformation with an Activation Tagging Construct

Mutants were generated using the activation tagging "ACTTAG" vector, pSKI015 (GI#6537289; Weigel D *et al.*, *supra*). Standard methods were used for the generation of *Arabidopsis* transgenic plants, and were essentially as described in published application PCT WO0183697. Briefly, T0 *Arabidopsis* (Col-0) plants were transformed with *Agrobacterium* carrying the pSKI015 vector, which comprises T-DNA derived from the *Agrobacterium* Ti plasmid, an herbicide resistance selectable marker gene, and the 4X CaMV 35S enhancer element. Transgenic plants were selected at the T1 generation based on herbicide resistance. T2 seed was collected from T1 plants and stored in an indexed collection, and a portion of the T2 seed was accessed for the screen.

Quantitative determination of seed fatty acid content was performed using the follows methods. An aliquot of 15 to 20 T2 seeds from each line tested, which generally

contained homozygous insertion, homozygous wild-type, and heterozygous genotypes in a standard 1:1:2 ratio, was massed on UMT-2 ultra-microbalance (Mettler-Toledo Co., Ohio, USA) and then transferred to a glass extraction vial. Whole seeds were transesterified in 500 ul 2.5% H<sub>2</sub>SO<sub>4</sub> in MeOH for 3 hours at 80 °C, following the method of

5 Browse et al. (Biochem J 235:25-31, 1986) with modifications. A known amount of heptadecanoic acid was included in the reaction as an internal standard. 750 ul of water and 400 ul of hexane were added to each vial, which was then shaken vigorously and allowed to phase separate. Reaction vials were loaded directly onto GC for analysis and the upper hexane phase was sampled by the autosampler. Gas chromatography with

10 Flame Ionization detection was used to separate and quantify the fatty acid methyl esters. Agilent 6890 Plus GC's were used for separation with Agilent Innowax columns (30m x 0.25mm ID, 250um film thickness). The carrier gas was Hydrogen at a constant flow of 2.5 ml/ minute. 1ul of sample was injected in splitless mode (inlet temperature 220°C, Purge flow 15ml/min at 1 minute). The oven was programmed for an initial temperature

15 of 105°C, initial time 0.5 minutes, followed by a ramp of 60°C per minute to 175°C, a 40°C /minute ramp to 260°C with a final hold time of 2 minutes. Detection was by Flame Ionization (Temperature 275°C, Fuel flow 30.0 ml/min, Oxidizer 400.0 ml/min). Instrument control and data collection and analysis was using the Millennium Chromatography Management System (Version 3.2, Waters Corporation, Milford, MA).

20 Integration and quantification were performed automatically, but all analyses were subsequently examined manually to verify correct peak identification and acceptable signal to noise ratio before inclusion of the derived results in the study.

The ACTTAG line designated W000086431 was identified as having a high oil phenotype. Specifically, oil constituted 34.8% of seed mass (w/w) compared to an

25 average oil content of 28.7% of other ACTTAG lines grown and analyzed in the same conditions (i.e. reference lines). Reanalysis of the same seed was performed in triplicate. Oil constituted 32.1% of seed mass, confirming an increase in oil content relative to reference.

The W000086431 line has three loci of T-DNA. One locus, was unlinked to the

30 phenotype, and two closely linked loci were identified which cosegregated with the high oil phenotype. T2 individuals homozygous for the high oil loci produced seed with an oil content of 115.4% of the reference, T2 individuals hemizygous for the loci produced seed with an oil content of 118.4% of the reference. T2 individuals lacking the HIO-30 locus had oil contents of 105% of the reference. Because the homozygotes and hemizygotes for

the high oil loci display a similar increase in oil content, it was determined that the high oil phenotype of W000068431 is dominant.

## EXAMPLE 2

### Characterization of the T-DNA Insertion in Plants Exhibiting the Altered Oil Content Phenotype.

We performed standard molecular analyses, essentially as described in patent application PCT WO0183697, to determine the site of the T-DNA insertion associated with the altered oil content phenotype. Briefly, genomic DNA was extracted from plants exhibiting the altered oil content phenotype. PCR, using primers specific to the pSKI015 vector, confirmed the presence of the 35S enhancer in plants from line W000086431, and Southern blot analysis verified the genomic integration of the ACTTAG T-DNA.

Plasmid rescue and/or inverse PCR were used to recover genomic DNA flanking the T-DNA insertion, which was then subjected to sequence analysis using a basic BLASTN search and/or a search of the Arabidopsis Information Resource (TAIR) database (available at the arabidopsis.org website). Seeds from 18 T3 families descended from the mutant were harvested. The seed oil content of these families was determined as described in Example 1. The genotypes of these families with respect to a T-DNA insert were determined by T-DNA specific PCR using primers that are specific to one of the corresponding genomic region. The average oil content of T3 families containing the T-DNA insert at locus 2 and 3 was higher than those families lacking the insert at the corresponding loci. Therefore, we concluded that locus 2 and/or 3 is linked to the high oil phenotype. By contrast, the average oil content of T3 families containing the T-DNA insert at locus 1 was lower than /approximately the same as those families lacking the insert at the corresponding locus, and we concluded that locus 1 is not linked to the high oil phenotype.

Sequence analysis revealed that the start codon of the nucleotide sequence presented as SEQ ID NO: 1, which we designated HIO30, was approximately 5.6 kb 5' of the upstream border of the T-DNA insert at locus 3.

## EXAMPLE 3

### Analysis of Arabidopsis HIO30 Sequence

Sequence analyses were performed with BLAST (Altschul *et al.*, 1997, J. Mol. Biol. 215:403-410), PFAM (Bateman *et al.*, 1999, Nucleic Acids Res 27:260-262),

PSORT (Nakai K, and Horton P, 1999, Trends Biochem Sci 24:34-6), and/or CLUSTAL (Thompson JD et al, 1994, Nucleic Acids Res 22:4673-4680).

BLASTN of SEQ ID NO:1 against GenBank sequences identified 6 Arabidopsis ESTs derived from cDNA clones derived from the same genomic region as At3g52260.

5 This gene appears to occur as a single copy in Arabidopsis.

BLASTN also identified additional ESTs from other plant species. The following three potato (*Solanum tuberosum*) ESTs were identified as candidate orthologs; BLAST scores are in parentheses: GI#s 13614224 (1054 4.2e-41), 18257466 (1031 4.8e-40), and 14266906 (1140 7.2e-50). Other plant ESTs were sorted by species, and then assembled  
10 into the least number of contigs, represented by SEQ ID NOs 3-6, and described further below. In most cases, the EST contigs represent partial coding regions. But, the entire cDNA sequence of the ortholog can be determined by someone skilled in molecular biology techniques.

SEQ ID NO:3 is from tomato (*Lycopersicon esculentum*), and is a contig of the  
15 following sequences: GI#s 5894121, 9503437, 5894459, and 5889410. It has 56% identity with SEQ ID NO:1.

SEQ ID NO:4 is from soy (*Glycine max*), and is a contig of GI#s 6913831, and 5760781. It has 66% identity with SEQ ID NO:1.

SEQ ID NO:5 is from corn (*Zea mays*), and is a contig of GI#s 21215230,  
20 5268759, and 22521540. It has 57% identity with SEQ ID NO:1.

SEQ ID NO:6 is from wheat (*Triticum aestivum*) and is a contig of GI#s 19955658 and 9364987. It has 62% identity with SEQ ID NO:1.

BLASTP analysis returned redundant entries for the At3g52260 gene product. The only other hits returned were numerous bacterial gene products corresponding to 23S  
25 RNA pseudouridylate synthase. No other plant sequences were returned.

PFam analysis detected an RNA pseudouridylate synthase domain (PF00849). In accordance with this functional prediction, PSORT2 predicts that the At3g52260 gene product is cytoplasmic (44%). The gene may have a regulatory function in the expression of genes involved in fatty acid metabolism or related pathways.

30

#### EXAMPLE 4

##### Confirmation of Phenotype/Genotype Association

RT-PCR analysis showed that the HIO30 gene was overexpressed in plants from the line displaying the HIO30 phenotype. Specifically, RNA was extracted from rosette leaves

and/or siliques of plants exhibiting the HIO30 phenotype collected at a variety of developmental stages and pooled. RT-PCR was performed using primers specific to the sequence presented as SEQ ID NO:1, to other predicted genes in the vicinity of the T-DNA insertion, and to a constitutively expressed actin gene (positive control). The results

5 showed that plants displaying the HIO30 phenotype over-expressed the mRNA for the HIO30 gene, indicating the enhanced expression of the HIO30 gene is correlated with the HIO30 phenotype.

The dominant inheritance pattern of the HIO30 phenotype is confirmed through genetic analysis. In general, genetic analysis involves the production and analysis of F1

10 hybrids. Typically, F1 crosses are carried out by collecting pollen from T2 plants, which is used to pollinate wild type plants. Such crosses are carried out by taking approximately 4 flowers from each selected individual plants, and using the T2 flower as the male pollen donor and flowers of the wild type plants as the female. 4-5 crosses are done for an individual of interest. Seed formed from crosses of the same individual are pooled,

15 planted and grown to maturity as F1 hybrids.

## IT IS CLAIMED:

1. A transgenic plant comprising a plant transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO30 polypeptide comprising the amino acid sequence of SEQ ID NO:2, or an ortholog thereof,  
5     whereby the transgenic plant has a high oil phenotype relative to control plants.
2. The transgenic plant of Claim 1, which is selected from the group consisting of rapeseed, soy, corn, sunflower, cotton, cocoa, safflower, oil palm, coconut palm, flax,  
10     castor and peanut.
3. A plant part obtained from the plant according to Claim 1.
4. The plant part of Claim 3, which is a seed.  
15
5. A method of producing oil comprising growing the transgenic plant of Claim 1 and recovering oil from said plant.
6. A method of producing a high oil phenotype in a plant, said method comprising:  
20     a) introducing into progenitor cells of the plant a plant transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO30 polypeptide comprising the amino acid sequence of SEQ ID NO:2, or an ortholog thereof, and  
       b) growing the transformed progenitor cells to produce a transgenic plant, wherein  
25     said polynucleotide sequence is expressed, and said transgenic plant exhibits an altered oil content phenotype relative to control plants.
7. A plant obtained by a method of Claim 6.
- 30   8. The plant of Claim 7, which is selected from the group consisting of rapeseed, soy, corn, sunflower, cotton, cocoa, safflower, oil palm, coconut palm, flax, castor and peanut.
9. A method of generating a plant having a high oil phenotype comprising identifying a plant that has an allele in its HIO30 gene that results in increased oil content compared to



plants lacking the allele and generating progeny of said identified plant, wherein the generated progeny inherit the allele and have the high oil phenotype.

10. The method of Claim 9 that employs candidate gene/QTL methodology.

5

11. The method of Claim 9 that employs TILLING methodology.

## SEQUENCE LISTING

&lt;110&gt; Agrinomics LLC

&lt;120&gt; GENERATION OF PLANTS WITH ALTERED OIL CONTENT

&lt;130&gt; AG03-076C-PC

&lt;150&gt; 60/434,763

&lt;151&gt; 2002-12-18

&lt;160&gt; 6

&lt;170&gt; PatentIn version 3.2

&lt;210&gt; 1

&lt;211&gt; 5034

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 1

```

atgccgcagg atcacgcttc gtgggatcgg aaagagctct tgaggcaaag gaaacacgat      60
aggcctgaac aatcttttga atccccgcct ttctgatgga gggattcgcc ttcttctcac      120
catgttcctc gagagttttc ttctcgtttg ggatctggag acttccgcag accttcttca      180
ttaacacagc tcttaagatt gataggaagt gaattgatga ggttactctt caaaggagca      240
atttttaata ctcaggggtg acggcaccag tttgtggagg agactagtca tggatacaca      300
tcttctcggt ccagtgcccg aatgtttgat aattataggc catcagcatc gcgtggagac      360
tgagatatata ccaggaattg cagggatgat agagtttctg taagccaaaa ggaatggaaa      420
tgcaatacat gggagatgag caacggatct tctagaagtt ttgagaggcc atttggtatt      480
agaaatggtc ggaggtcagt tgatgaaagg ccgctacatg cttcagatac tcattctacc      540
gtggtgaact ctttggatcc agccaactcg gctcattatc tggacaatga gatcagtacc      600
ccagtacggt ctcttaaaat taaaaatgag cataaatttt cagatcaaag gttatcactt      660
ccttcagatc ctcattctga atgtattagc ttgtttgaac ggccttcttc tgagaacaat      720
tatggcaata aggtttgttc accagcaaag caatgcaatg atttgatgta tggtcgaagg      780
ttagttagtg ataattcatt agatgctcca atccccaatg cagagctgga ggggacttgg      840
gaacaacttc gcctgaaaga cccgcaagat aacaatagtt tacatgggat caatgatata      900
gacggtgata ggaaatgtgc aaaggagagt tctctgggag caactgggaa acttccactg      960
tggaatagtt ctgggagttt tgcattctcag agttcagggt ttagtcattc aagtagcttg     1020
aaaagcttgg gggctgttga ttccagcgat cggaagattg aggttcttcc taaaattggt     1080
actgtgactc aatcttcttc aggagacgct actgcctgtg ccacaactac tcatctttct     1140
gaggagatga gttctagaaa gaaacaacgt ctcgggtggg gtgagggact ggcgaaatat     1200
gagaaaaaga aagttgatgt taaccctaat gaagatggaa caacattgat ggaaaacggt     1260

```

ttagaggaac	tacattcggt	aaacaaaaat	attgctgata	aaagtccac	agcagccatt	1320
gttccagatt	atgggtcccc	tacaacacca	tcctctgtag	cttgcagttc	atcaccaggg	1380
tttgctgata	aatcatctcc	gaaggctgct	atagctgcta	gtgatgtcag	taacatgtgc	1440
cgttcgccta	gtcccgtgtc	tagtattcac	cttgaacgat	tcccaatcaa	tatcgaggag	1500
ctcgataaca	tctcaatgga	gcgttttggc	tgtttactca	atgagttact	tggtactgat	1560
gattctggta	caggggattc	cagttctgtc	caattgacat	caatgaacac	attacttgcc	1620
tggaaaggtg	aaattttgaa	agctgtggag	atgactgaat	cagaaattga	tctccttgaa	1680
aacaaacata	ggacactaaa	gcttgaaggt	agaagacact	ctcgtgttgt	tggaccaggt	1740
tcatactggt	gtgatggaga	tgcaaagtgt	ccaaggagc	aggcttcttg	tagtttggat	1800
cctaaggcaa	cagcttcttc	tgtagctaaa	acactgggtga	gagctcctgt	gcatcaggct	1860
ggtttagcca	aggttcctgc	tgatgttttt	gaagatagtc	ctgggggaagt	taaacctcta	1920
tcccaatctt	ttgccactgt	tgaaagagag	gaagatatac	tgcccatacc	atctatgaag	1980
gcagctgttt	cttcgaaaga	gattaacaca	cctgcttttg	ccaatcagga	aactattgag	2040
gtttcttctg	ctgatgacag	catggcctcc	aaagaagact	tgttctgggc	taagttatta	2100
tctgccataa	agaaatatgc	ttgtgaatca	tctggagtat	tcaatcaatt	gcttccaaga	2160
gattttaatt	cgtctgacaa	ctcaagattc	cctggcatat	gtcaaacgca	gtttgattct	2220
catgtccaag	aaaaaattgc	agatagggta	ggcctattga	gagctagggg	gaaaatttta	2280
ctccttcagt	ttaaagcggt	tcagctctca	tggagaagag	atttggtatca	gctagcttta	2340
gcaaagtacc	aatcaaagtc	tagcaaaaaa	acagaactat	atccgaatgc	aaaaaatgga	2400
gggtatctga	agcttcccca	atctgtacgc	ctgaggttct	cttcttcagc	tccaagaagg	2460
gatagtgtag	tccccacaac	agagctcgta	agttatatgg	aaaagctact	tccgggtacc	2520
catctaaagc	cttttagaga	cattttgaaa	atgcttgcta	tgattttgga	tgagaaagag	2580
aggggtgatgt	cgaggtttat	ttctagcaat	ggactgattg	aagatccatg	tgacgttgag	2640
aaggaaagaa	caatgattaa	tccttggacc	tcagaggaga	aagaaatctt	tctgaatttg	2700
ctagcaatgc	atgggaagga	tttcaagaag	attgcttcat	ctcttacc	aaagacaact	2760
gcggactgta	ttgattacta	ctacaaaaac	cacaagtctg	attgttttgg	gaaaataaag	2820
aagcagcgtg	cttatggtaa	ggaagggaag	cacacctaca	tgttggctcc	acgaaaaaag	2880
tggaaacgtg	agatgggggc	tgctctctct	gatatttttag	gggatgtctc	cattatagca	2940
gcaaacgctg	gaaagggtgc	atcaaccagg	ccgatctctt	ccaaaaagat	cacccttaga	3000
ggttgcagca	gtgctaattc	attgcagcac	gatggaaata	actctgaagg	gtgctcctac	3060
agttttgatt	tcccacgtaa	gagaactgct	ggtgcagatg	ttttagctgt	tggctccttg	3120

tcaccagagc agataaatc ttgcttaagg acttctgtga gctctagaga gaggtgtatg 3180  
gatcatctga agtttaatca tgtcgtaaag aaacctcgga tatctcatac tctacataat 3240  
gagaacagca atactctaca caatgagaac agcaacgaag aagatgactc atgttcggaa 3300  
gagagctgtg gggaaacagg tcctattcac tggacagatg atgagagatc tgcctttata 3360  
caggggtttt cgctttttgg caagaatttt gcttcaatat caaggtagct cgggacaaga 3420  
tctccagatc agtgtaaggt tttcttcagc aaagttcgga aatgtcttgg gttggaatct 3480  
ataaagtttg gatctggaaa tgtaagcaca tccgtaagtg ttgataatgg caatgagggg 3540  
gggtgggagcg acttggaaga tccttgtcct atggagagta actctggcat agtgaataat 3600  
ggagtttgtg ccaagatggg tatgaattct cctacctcac cttttaatat gaatcaggat 3660  
gggtgttaatc aatcaggctc tgcaaagtgt aaagccgacc ttagtagatc agaagaagag 3720  
aatgggcaga aatatttgtg tctgaaagat gataataatc tcgtgaacaa tgcataatgtc 3780  
aatggcggtt tcccagatct agtttcagaa tcttgtagag atttggtaga tattaatact 3840  
gttgagagcc agtctcaggc tgccggaaaa agcaagagca atgatctcat gtcaatggaa 3900  
atcgatgaag gtgtcttaac atctgtcact atatcttccg agccattgta ttgtggccta 3960  
agtgttcttt ccaatgttat tgtggaaacc cctacagaaa tctcacgaaa gggctcagga 4020  
gatcaagggtg ctacaatgcc taaatttagt tcaaagaatc aagatggagt gatgcaagct 4080  
gcaaacagaa ccagaaattc tggccttgaa cctgaaagtg caccttcagg ttccaggtac 4140  
cctgagtgtc ttcacatgt tccgattgag gtgtgtacgg aaaaccctat aggcgtcagt 4200  
gcaccacgag gaaatccaaa ttgccatgca gagtccgagt caggaaattc tcttgttgga 4260  
caagttgacg aaacacatga cttgggttgg cccaagaaca atctggaatt ggatgggagg 4320  
cttcagggtt taggccatgt aaacctgag cagattgggtc tactaaaagc gaccaatata 4380  
gaatcttgtc aaaatcccca gagatcagtc acccaagatc tgagcaggat aagtagatca 4440  
aaatctgatt tgatcgtaaa aaccaacgt acaggtgaag gcttctcact caccaagtgt 4500  
actagttcag ctcttaagcc tctggcagta tcccataaag agggcagatc tggatcatagc 4560  
aggagccatt cgttttagtt gtctgatact gagagactcc acaagaatgg agatgtgaaa 4620  
ctgtttggta cagtacttac tactgatgag aatggaataa aacaaaaaca caatccatgt 4680  
ggaattgtca ggtcatcatc aaccttgagc agggaccatg atacaagaca tcattacatt 4740  
aatcagcaac accttcagaa cgttcccatt acgagctacg gtttttggga tggcaacaga 4800  
attcaaaccg ggctcacatc tttgccagag tcggccaagt tgcttgcaag ttgccctgaa 4860  
gcattttcca cgcactctaaa gcagcaagtt ggtaacagca aagagattct ggtggatgtt 4920  
aatgggtggaa ttttgagctt tggttaagcat aacgaagata gagctgagtc ctcaagcgct 4980

aaggatgaag gtaacatagg aggggtaaataa ggtgtagcag aggcagccac gtga 5034

<210> 2  
 <211> 1677  
 <212> PRT  
 <213> Arabidopsis thaliana  
 <400> 2

Met Pro Gln Asp His Ala Ser Trp Asp Arg Lys Glu Leu Leu Arg Gln  
 1 5 10 15

Arg Lys His Asp Arg Pro Glu Gln Ser Phe Glu Ser Pro Pro Phe Arg  
 20 25 30

Trp Arg Asp Ser Pro Ser Ser His His Val Pro Arg Glu Phe Ser Ser  
 35 40 45

Arg Leu Gly Ser Gly Asp Phe Arg Arg Pro Ser Ser Leu Thr Gln Leu  
 50 55 60

Leu Arg Leu Ile Gly Ser Glu Leu Met Arg Leu Leu Phe Lys Gly Ala  
 65 70 75 80

Ile Phe Asn Thr Gln Gly Gly Arg His Gln Phe Val Glu Glu Thr Ser  
 85 90 95

His Gly Tyr Thr Ser Ser Arg Ser Ser Ala Arg Met Phe Asp Asn Tyr  
 100 105 110

Arg Pro Ser Ala Ser Arg Gly Asp Trp Arg Tyr Thr Arg Asn Cys Arg  
 115 120 125

Asp Asp Arg Val Ser Val Ser Gln Lys Glu Trp Lys Cys Asn Thr Trp  
 130 135 140

Glu Met Ser Asn Gly Ser Ser Arg Ser Phe Glu Arg Pro Phe Gly Ile  
 145 150 155 160

Arg Asn Gly Arg Arg Ser Val Asp Glu Arg Pro Leu His Ala Ser Asp  
 165 170 175

Thr His Ser Thr Val Val Asn Ser Leu Asp Pro Ala Asn Ser Ala His  
 180 185 190

Tyr Leu Asp Asn Glu Ile Ser Thr Pro Val Arg Ser Leu Lys Ile Lys  
 195 200 205

Asn Glu His Lys Phe Ser Asp Gln Arg Leu Ser Leu Pro Ser Asp Pro  
 210 215 220

His Ser Glu Cys Ile Ser Leu Phe Glu Arg Pro Ser Ser Glu Asn Asn  
 225 230 235 240

Tyr Gly Asn Lys Val Cys Ser Pro Ala Lys Gln Cys Asn Asp Leu Met  
 245 250 255

Tyr Gly Arg Arg Leu Val Ser Asp Asn Ser Leu Asp Ala Pro Ile Pro  
 260 265 270

Asn Ala Glu Leu Glu Gly Thr Trp Glu Gln Leu Arg Leu Lys Asp Pro  
 275 280 285

Gln Asp Asn Asn Ser Leu His Gly Ile Asn Asp Ile Asp Gly Asp Arg  
 290 295 300

Lys Cys Ala Lys Glu Ser Ser Leu Gly Ala Thr Gly Lys Leu Pro Leu  
 305 310 315 320

Trp Asn Ser Ser Gly Ser Phe Ala Ser Gln Ser Ser Gly Phe Ser His  
 325 330 335

Ser Ser Ser Leu Lys Ser Leu Gly Ala Val Asp Ser Ser Asp Arg Lys  
 340 345 350

Ile Glu Val Leu Pro Lys Ile Val Thr Val Thr Gln Ser Ser Ser Gly  
 355 360 365

Asp Ala Thr Ala Cys Ala Thr Thr Thr His Leu Ser Glu Glu Met Ser  
 370 375 380

Ser Arg Lys Lys Gln Arg Leu Gly Trp Gly Glu Gly Leu Ala Lys Tyr  
 385 390 395 400

Glu Lys Lys Lys Val Asp Val Asn Pro Asn Glu Asp Gly Thr Thr Leu  
 405 410 415

Met Glu Asn Gly Leu Glu Glu Leu His Ser Leu Asn Lys Asn Ile Ala  
 420 425 430

Asp Lys Ser Pro Thr Ala Ala Ile Val Pro Asp Tyr Gly Ser Pro Thr  
 435 440 445

Thr Pro Ser Ser Val Ala Cys Ser Ser Ser Pro Gly Phe Ala Asp Lys

450		455		460	
Ser Ser Pro Lys Ala Ala Ile Ala Ala Ser Asp Val Ser Asn Met Cys					
465		470		475	480
Arg Ser Pro Ser Pro Val Ser Ser Ile His Leu Glu Arg Phe Pro Ile					
	485		490		495
Asn Ile Glu Glu Leu Asp Asn Ile Ser Met Glu Arg Phe Gly Cys Leu					
	500		505		510
Leu Asn Glu Leu Leu Gly Thr Asp Asp Ser Gly Thr Gly Asp Ser Ser					
	515		520		525
Ser Val Gln Leu Thr Ser Met Asn Thr Leu Leu Ala Trp Lys Gly Glu					
	530		535		540
Ile Leu Lys Ala Val Glu Met Thr Glu Ser Glu Ile Asp Leu Leu Glu					
	545		550		555
					560
Asn Lys His Arg Thr Leu Lys Leu Glu Gly Arg Arg His Ser Arg Val					
	565		570		575
Val Gly Pro Ser Ser Tyr Cys Cys Asp Gly Asp Ala Asn Val Pro Lys					
	580		585		590
Glu Gln Ala Ser Cys Ser Leu Asp Pro Lys Ala Thr Ala Ser Ser Val					
	595		600		605
Ala Lys Thr Leu Val Arg Ala Pro Val His Gln Ala Gly Leu Ala Lys					
	610		615		620
Val Pro Ala Asp Val Phe Glu Asp Ser Pro Gly Glu Val Lys Pro Leu					
	625		630		635
					640
Ser Gln Ser Phe Ala Thr Val Glu Arg Glu Glu Asp Ile Leu Pro Ile					
	645		650		655
Pro Ser Met Lys Ala Ala Val Ser Ser Lys Glu Ile Asn Thr Pro Ala					
	660		665		670
Phe Ala Asn Gln Glu Thr Ile Glu Val Ser Ser Ala Asp Asp Ser Met					
	675		680		685
Ala Ser Lys Glu Asp Leu Phe Trp Ala Lys Leu Leu Ser Ala Asn Lys					
	690		695		700

Lys Tyr Ala Cys Glu Ser Ser Gly Val Phe Asn Gln Leu Leu Pro Arg  
 705 710 715 720  
 Asp Phe Asn Ser Ser Asp Asn Ser Arg Phe Pro Gly Ile Cys Gln Thr  
 725 730 735  
 Gln Phe Asp Ser His Val Gln Glu Lys Ile Ala Asp Arg Val Gly Leu  
 740 745 750  
 Leu Arg Ala Arg Glu Lys Ile Leu Leu Leu Gln Phe Lys Ala Phe Gln  
 755 760 765  
 Leu Ser Trp Lys Lys Asp Leu Asp Gln Leu Ala Leu Ala Lys Tyr Gln  
 770 775 780  
 Ser Lys Ser Ser Lys Lys Thr Glu Leu Tyr Pro Asn Ala Lys Asn Gly  
 785 790 795 800  
 Gly Tyr Leu Lys Leu Pro Gln Ser Val Arg Leu Arg Phe Ser Ser Ser  
 805 810 815  
 Ala Pro Arg Arg Asp Ser Val Val Pro Thr Thr Glu Leu Val Ser Tyr  
 820 825 830  
 Met Glu Lys Leu Leu Pro Gly Thr His Leu Lys Pro Phe Arg Asp Ile  
 835 840 845  
 Leu Lys Met Pro Ala Met Ile Leu Asp Glu Lys Glu Arg Val Met Ser  
 850 855 860  
 Arg Phe Ile Ser Ser Asn Gly Leu Ile Glu Asp Pro Cys Asp Val Glu  
 865 870 875 880  
 Lys Glu Arg Thr Met Ile Asn Pro Trp Thr Ser Glu Glu Lys Glu Ile  
 885 890 895  
 Phe Leu Asn Leu Leu Ala Met His Gly Lys Asp Phe Lys Lys Ile Ala  
 900 905 910  
 Ser Ser Leu Thr Gln Lys Thr Thr Ala Asp Cys Ile Asp Tyr Tyr Tyr  
 915 920 925  
 Lys Asn His Lys Ser Asp Cys Phe Gly Lys Ile Lys Lys Gln Arg Ala  
 930 935 940  
 Tyr Gly Lys Glu Gly Lys His Thr Tyr Met Leu Ala Pro Arg Lys Lys



945				950				955				960			
Trp	Lys	Arg	Glu	Met	Gly	Ala	Ala	Ser	Leu	Asp	Ile	Leu	Gly	Asp	Val
			965						970				975		
Ser	Ile	Ile	Ala	Ala	Asn	Ala	Gly	Lys	Val	Ala	Ser	Thr	Arg	Pro	Ile
			980					985					990		
Ser	Ser	Lys	Lys	Ile	Thr	Leu	Arg	Gly	Cys	Ser	Ser	Ala	Asn	Ser	Leu
		995					1000					1005			
Gln	His	Asp	Gly	Asn	Asn	Ser	Glu	Gly	Cys	Ser	Tyr	Ser	Phe	Asp	
	1010					1015					1020				
Phe	Pro	Arg	Lys	Arg	Thr	Ala	Gly	Ala	Asp	Val	Leu	Ala	Val	Gly	
	1025					1030					1035				
Pro	Leu	Ser	Pro	Glu	Gln	Ile	Asn	Ser	Cys	Leu	Arg	Thr	Ser	Val	
	1040					1045					1050				
Ser	Ser	Arg	Glu	Arg	Cys	Met	Asp	His	Leu	Lys	Phe	Asn	His	Val	
	1055					1060					1065				
Val	Lys	Lys	Pro	Arg	Ile	Ser	His	Thr	Leu	His	Asn	Glu	Asn	Ser	
	1070					1075					1080				
Asn	Thr	Leu	His	Asn	Glu	Asn	Ser	Asn	Glu	Glu	Asp	Asp	Ser	Cys	
	1085					1090					1095				
Ser	Glu	Glu	Ser	Cys	Gly	Glu	Thr	Gly	Pro	Ile	His	Trp	Thr	Asp	
	1100					1105					1110				
Asp	Glu	Arg	Ser	Ala	Phe	Ile	Gln	Gly	Phe	Ser	Leu	Phe	Gly	Lys	
	1115					1120					1125				
Asn	Phe	Ala	Ser	Ile	Ser	Arg	Tyr	Val	Gly	Thr	Arg	Ser	Pro	Asp	
	1130					1135					1140				
Gln	Cys	Lys	Val	Phe	Phe	Ser	Lys	Val	Arg	Lys	Cys	Leu	Gly	Leu	
	1145					1150					1155				
Glu	Ser	Ile	Lys	Phe	Gly	Ser	Gly	Asn	Val	Ser	Thr	Ser	Val	Ser	
	1160					1165					1170				
Val	Asp	Asn	Gly	Asn	Glu	Gly	Gly	Gly	Ser	Asp	Leu	Glu	Asp	Pro	
	1175					1180					1185				

Cys	Pro	Met	Glu	Ser	Asn	Ser	Gly	Ile	Val	Asn	Asn	Gly	Val	Cys
1190						1195					1200			
Ala	Lys	Met	Gly	Met	Asn	Ser	Pro	Thr	Ser	Pro	Phe	Asn	Met	Asn
1205						1210					1215			
Gln	Asp	Gly	Val	Asn	Gln	Ser	Gly	Ser	Ala	Asn	Val	Lys	Ala	Asp
1220						1225					1230			
Leu	Ser	Arg	Ser	Glu	Glu	Glu	Asn	Gly	Gln	Lys	Tyr	Leu	Cys	Leu
1235						1240					1245			
Lys	Asp	Asp	Asn	Asn	Leu	Val	Asn	Asn	Ala	Tyr	Val	Asn	Gly	Gly
1250						1255					1260			
Phe	Pro	Ser	Leu	Val	Ser	Glu	Ser	Cys	Arg	Asp	Leu	Val	Asp	Ile
1265						1270					1275			
Asn	Thr	Val	Glu	Ser	Gln	Ser	Gln	Ala	Ala	Gly	Lys	Ser	Lys	Ser
1280						1285					1290			
Asn	Asp	Leu	Met	Ser	Met	Glu	Ile	Asp	Glu	Gly	Val	Leu	Thr	Ser
1295						1300					1305			
Val	Thr	Ile	Ser	Ser	Glu	Pro	Leu	Tyr	Cys	Gly	Leu	Ser	Val	Leu
1310						1315					1320			
Ser	Asn	Val	Ile	Val	Glu	Thr	Pro	Thr	Glu	Ile	Ser	Arg	Lys	Gly
1325						1330					1335			
Ser	Gly	Asp	Gln	Gly	Ala	Thr	Met	Pro	Lys	Phe	Ser	Ser	Lys	Asn
1340						1345					1350			
Gln	Asp	Gly	Val	Met	Gln	Ala	Ala	Asn	Arg	Thr	Arg	Asn	Ser	Gly
1355						1360					1365			
Leu	Glu	Pro	Glu	Ser	Ala	Pro	Ser	Gly	Phe	Arg	Tyr	Pro	Glu	Cys
1370						1375					1380			
Leu	His	His	Val	Pro	Ile	Glu	Val	Cys	Thr	Glu	Asn	Pro	Ile	Gly
1385						1390					1395			
Val	Ser	Ala	Pro	Arg	Gly	Asn	Pro	Asn	Cys	His	Ala	Glu	Ser	Glu
1400						1405					1410			
Ser	Gly	Asn	Ser	Leu	Val	Gly	Gln	Val	Asp	Glu	Thr	His	Asp	Leu

1415		1420		1425
Gly Trp Pro Lys Asn Asn Leu Glu Leu Asp Gly Arg Leu Gln Val				
1430		1435		1440
Leu Gly His Val Asn Pro Glu Gln Ile Gly Leu Leu Lys Ala Thr				
1445		1450		1455
Asn Thr Glu Ser Cys Gln Asn Pro Gln Arg Ser Val Thr Gln Asp				
1460		1465		1470
Leu Ser Arg Ile Ser Arg Ser Lys Ser Asp Leu Ile Val Lys Thr				
1475		1480		1485
Gln Arg Thr Gly Glu Gly Phe Ser Leu Thr Lys Cys Thr Ser Ser				
1490		1495		1500
Ala Pro Lys Pro Leu Ala Val Ser His Lys Glu Gly Arg Ser Gly				
1505		1510		1515
His Ser Arg Ser His Ser Phe Ser Leu Ser Asp Thr Glu Arg Leu				
1520		1525		1530
His Lys Asn Gly Asp Val Lys Leu Phe Gly Thr Val Leu Thr Thr				
1535		1540		1545
Asp Glu Asn Gly Ile Lys Gln Lys His Asn Pro Cys Gly Ile Val				
1550		1555		1560
Arg Ser Ser Ser Thr Leu Ser Arg Asp His Asp Thr Arg His His				
1565		1570		1575
Tyr Ile Asn Gln Gln His Leu Gln Asn Val Pro Ile Thr Ser Tyr				
1580		1585		1590
Gly Phe Trp Asp Gly Asn Arg Ile Gln Thr Gly Leu Thr Ser Leu				
1595		1600		1605
Pro Glu Ser Ala Lys Leu Leu Ala Ser Cys Pro Glu Ala Phe Ser				
1610		1615		1620
Thr His Leu Lys Gln Gln Val Gly Asn Ser Lys Glu Ile Leu Val				
1625		1630		1635
Asp Val Asn Gly Gly Ile Leu Ser Phe Gly Lys His Asn Glu Asp				
1640		1645		1650

Arg Ala Glu Ser Ser Ser Ala Lys Asp Glu Gly Asn Ile Gly Gly  
 1655 1660 1665

Val Asn Gly Val Ala Glu Ala Ala Thr  
 1670 1675

<210> 3  
 <211> 1012  
 <212> DNA  
 <213> *Lycopersicon esculentum*

<400> 3  
 caggaaaattt gtagaatttg aatttgagtt taatatatttg gccagaaaatt tgttgatttc 60  
 ttcaagtttt ggattaatct gctgctgatt gtttcaggaa gttgcttctg gcgatattcc 120  
 ctactccgag ttacgaatgt cagaaaaagc agagcttgca gcggtgaggg cgtactttgg 180  
 agtgctgtgg ccgcaacgta atgagggatt atcctaccat gacatcgccc gaccaccga 240  
 tgctgggtctt acattgatcg aattctactt taggaagtag aaaaattcag ctccctttaca 300  
 aggttggttg cagagaattc aaaataaaca gataacaatt gatggtaaag ttgttatctt 360  
 accagatact gaactcagag cagggtgctga attagtatat catcgccctt cttggagaga 420  
 acctgatgca ccttacttgc tagaagtact atttgaagat gactacttga ttgttgtaaa 480  
 taaaccttct ggtttgcaag ttcttcctgg agggttatat cagcagcgga ccgtcttgac 540  
 gcaactccag tggcatgcat gtaagctgac aaccacttcg tcaggttgtc aaaaaacaca 600  
 tccagtccca gttcatcgct taggaagggg tacatcagga atactgctct gtgcaaaaac 660  
 aaagctttgt aaatctcgcc ttgcagcata ttttgctgag gggacgtcag ttgttggaaga 720  
 aaaatgcacc aactcagagt gcaatacaat gaggaagatt tgcaagatat atcgggcgct 780  
 agtaagtggg gtgatggata tggatgaggg tgtcatcaag caaccaattg gtacaattaa 840  
 atatcctgga gttgctaaag ggttgatgtg tgcttctcct tcagggaagc cagctttgag 900  
 cagtgttcgc gttcttgaaa gagattcaga gagtaactgc acattgggtc aggttgaaat 960  
 tcaatctgga aggccacacc aaatccgcat ccacctctct ttcataggat at 1012

<210> 4  
 <211> 737  
 <212> DNA  
 <213> *Glycine max*

<400> 4  
 cttggcctga atgcaacgac ggtctgtcct acgacgacgt cgttcgagcc tctgatgccg 60  
 gggcgacact catagagttt tactccacca agtacaagag ttctgctccc ttacaagggt 120  
 ggttgacgag aataaaaagt gggcagataa cagttgatgg aggagttggt actgattcta 180

```

acacagtcct cagagttgga tcaaagctaa tctatcatag acttccatgg aaggagccag      240
atgcaccgca catgatcgac gtcttatatg aagatgatga catgattgct ctaaataaac      300
cgtctggcct gcaagttttg cctggaggtc tctaccagca gaggacaatt ttaacacagc      360
ttcaatggga agccaacaat caggggtacct gtgaaatgca caaaaggctg cattctggtc      420
ccgtgcatcg cctagggagg gggacttcag gaattttatt atgtgcgaag aaaaaactag      480
ccagagctcg tcttgcatct cattttgctg acggaacttc tcacgttgga ggaaaaagag      540
atacaaagca ggaacttggg aagattgcaa agatgtaccg agctcttggt agtgggatag      600
ttgagaatga caaggtgact attaatcaac caattggaat agtaaaatat cctggtgttg      660
ctaaaggggt atacgttgct tctgaatcag gaaaaccagc actcagtgtg gtggacattc      720
tagagacgaa catacaa                                     737

```

```

<210> 5
<211> 1308
<212> DNA
<213> Zea mays

```

```

<220>
<221> misc_feature
<222> (661)..(665)
<223> n is a, c, g, or t

```

```

<220>
<221> misc_feature
<222> (1186)..(1186)
<223> n is a, c, g, or t

```

```

<400> 5
gcacgaggcg gccgcggcag aaacacagat gggcgacggg gctccgccgc caggggctct      60
atactccttc ggaacgccgt ggccggagct caatcaaggc ctcacctaca ccgacacgtt      120
ccgttgcgct gatgcggacg ccgctaccac cttgattgag ttctactcta ctaaccacaa      180
gagctcggcg ccattgccag ggtggatcaa gaggattcgt aatgggcaga taaccgttga      240
tggtgaagtt gtcactgatc cagatatgac tctgggtggat ggggtctaagt tggatatatca      300
tcgttttcct tggcaggagc catttgccgc gtatttgctg gaagtgctct acgaggatga      360
tgacatgggt gcccttaata agccttctgg cttgcaagtt ctgcctaaag gactgtttca      420
gcagcgaact gtttttagcac agcttcaatt gaaagactgg aagatggcct catcttgccg      480
gttcaagaga aaagatgtgc agtcacatcc agtacctgtt catcgtttag ggaggggaac      540
atcaggcctc ctgctttgtg ccaagacaaa agttgccaaa gttcgacttg catcttattt      600
tgctgaaggt gctataaatg ctgcaaagaa aagggataaa tcagagttca gtgaagagcg      660
nnnnntttca aaattttatc gagccttagt gactggcata cttgatgatg atgaggttgt      720

```

```

tgttacgcaa cctatagggg tagttcatta tcctggaggt gcagagggac tttatgcagc      780
atgttcctca ggaaagccag caatgagcaa agtatgtggt cttgagagac ttgcacacca      840
aaatcacaca ctggtccagg ttgaaattca ttcaggacga cctcaccaaa taaggatata      900
ccttgcatac attgggcacc cacttgtaga tgaccctctc tatgggtattg gtgggcaccc      960
caattttgtt gagccagaat ctactggcac agatagttct tttgcatctg atgggggtta     1020
tgagagacct ttgcaacctg ttcttgaga ctgtgggtat cacctgcatg cacattggct     1080
ggttctttgc caccacaaca ccaataagat ggtaaaaatt accgctcctc ttccacaaat     1140
tctacagaca cgggaggaac gccgcgctgc agctgagcaa accgngggtt gaacatgtag     1200
aatcttgaaa atgtatattt cttgaagtta gcaagcagca ggttctcaca gacgttagag     1260
ttagacactc agacatctgc tcctctgtca actgtacaac ggcgagct                     1308

```

```

<210> 6
<211> 566
<212> DNA
<213> Triticum aestivum

```

```

<400> 6
atactccgcc gctgcaggcg aactacgtct tcgggagggc atggccggat ctcaacgaag      60
gactctccta caccgatacg ttccgcggcg ctgatgcgga aaccaccgcc accttgacca     120
atttctactc tgagaactac aagagctcgg cgccattgcc aggggtggatt cataggattc     180
gcaatggaca gataacggtt gatggccaag ttgtcactga tccagatatg attctcaggg     240
agggttctaa gttggtatat catgcctcg catggaagga gccatttgca ccacatttgc     300
ttcaagtgtt ttatgaagat gacgacatgg tagcccttaa taagccttcc ggtttgcaag     360
ttctgcaaaa aggactcttc cagcagcgca ctgttctagc acaacttcag tggaaagagt     420
ggaagatgcc cccatcaagc tgctctaaga gaaaaaatgt gcagttacat cctgtacctg     480
ttcatcgatt aggaaggggc acgtcaggtc tactgctttg tgccaagaca aagcttgcca     540
aagttcaact tgcattcttat tttgca                                         566

```